

CONTINUATION APPLICATION  
FOR  
UNITED STATES LETTERS PATENT

TITLE: FATTY ACID DESATURASES AND MUTANT  
SEQUENCES THEREOF

APPLICANT: LORIN R. DEBONTE, PH.D., ZHEGONG FAN AND GUO-  
HUA MIAO

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL639959554US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

1-29-01  
Date of Deposit

  
Signature

Vince Defante

Typed or Printed Name of Person Signing Certificate

07'48-063003-012901

FATTY ACID DESATURASES AND MUTANT SEQUENCES THEREOF

5

Cross-Reference to Related Application

This application is a continuation-in-part of PCT Application Serial No. PCT/US96/20090, filed December 13, 1996, which is a continuation-in-part of U.S. Application Serial No. 08/572,027, filed December 14, 1995.

10

Technical Field

This invention relates to fatty acid desaturases and nucleic acids encoding desaturase proteins. More particularly, the invention relates to nucleic acids encoding delta-12 and delta-15 fatty acid desaturase proteins that affect fatty acid composition in plants, polypeptides produced from such nucleic acids and plants expressing such nucleic acids.

Background of the Invention

Many breeding studies have been conducted to improve the fatty acid profile of Brassica varieties. Pleines and Freidt, Fat Sci. Technol., 90(5), 167-171 (1988) describe plant lines with reduced  $C_{18:3}$  levels (2.5-5.8%) combined with high oleic content (73-79%). Rakow and McGregor, J. Amer. Oil Chem. Soc., 50, 400-403 (Oct. 1973) discuss problems associated with selecting mutants for linoleic and linolenic acids. In, Can. J. Plant Sci., 68, 509-511 (Apr. 1988) Stellar summer rape producing seed oil with 3% linolenic acid and 28% linoleic acid is disclosed. Roy and Tarr, Z. Pflanzenzuchtg, 95(3), 201-209 (1985) teaches transfer of genes through an interspecific cross from Brassica juncea into Brassica napus resulting in a reconstituted line combining high linoleic with low linolenic acid content. Roy and Tarr, Plant Breeding, 98,

89-96 (1987) discuss prospects for development of *B. napus* L. having improved linolenic and linolenic acid content. European Patent application 323,753 published July 12, 1989 discloses seeds and oils having greater than 79% oleic acid combined with less than 3.5% linolenic acid. Canvin, Can. J. Botany, 43, 63-69 (1965) discusses the effect of temperature on the fatty acid composition of oils from several seed crops including rapeseed.

Mutations typically are induced with extremely high doses of radiation and/or chemical mutagens (Gaul, H. Radiation Botany (1964) 4:155-232). High dose levels which exceed LD50, and typically reach LD90, led to maximum achievable mutation rates. In mutation breeding of *Brassica* varieties high levels of chemical mutagens alone or combined with radiation have induced a limited number of fatty acid mutations (Rakow, G.Z. Pflanzenzuchtg (1973) 69:62-82). The low  $\alpha$ -linolenic acid mutation derived from the Rakow mutation breeding program did not have direct commercial application because of low seed yield. The first commercial cultivar using the low  $\alpha$ -linolenic acid mutation derived in 1973 was released in 1988 as the variety Stellar (Scarth, R. et al., Can. J. Plant Sci. (1988) 68:509-511). Stellar was 20% lower yielding than commercial cultivars at the time of its release.

Alterations in fatty acid composition of vegetable oils is desirable for meeting specific food and industrial uses. For example, *Brassica* varieties with increased monounsaturate levels (oleic acid) in the seed oil, and products derived from such oil, would improve lipid nutrition. Canola lines which are low in polyunsaturated fatty acids and high in oleic acid tend to have higher oxidative stability, which is a useful trait for the retail food industry.

Delta-12 fatty acid desaturase (also known as oleic desaturase) is involved in the enzymatic conversion of oleic acid to linoleic acid. Delta-15 fatty acid desaturase (also known as linoleic acid desaturase) is involved in the enzymatic conversion of linoleic acid to  $\alpha$ -linolenic acid. A microsomal delta-12 desaturase has been cloned and characterized using T-DNA tagging. Okuley, et al., Plant Cell 6:147-158 (1994). The nucleotide sequences of higher plant genes encoding microsomal delta-12 fatty acid desaturase are described in Lightner et al., WO94/11516. Sequences of higher plant genes encoding microsomal and plastid delta-15 fatty acid desaturases are disclosed in Yadav, N., et al., Plant Physiol., 103:467-476 (1993), WO 93/11245 and Arondel, V. et al., Science, 258:1353-1355 (1992). However, there are no teachings that disclose mutations in delta-12 or delta-15 fatty acid desaturase coding sequences from plants. There is a need in the art for more efficient methods to develop plant lines that contain delta-12 or delta-15 fatty acid desaturase gene sequence mutations effective for altering the fatty acid composition of seeds.

#### Summary of the Invention

The invention comprises *Brassicaceae* or *Helianthus* seeds, plants and plant lines having at least one mutation that controls the levels of unsaturated fatty acids in plants. One embodiment of the invention is an isolated nucleic acid fragment comprising a nucleotide sequence encoding a mutation from a mutant delta-12 fatty acid desaturase conferring altered fatty composition in seeds when the fragment is present in a plant. A preferred sequence comprises a mutant sequence as shown in Fig. 2. Another embodiment of the invention is an isolated nucleic

acid fragment comprising a nucleotide sequence encoding a mutation from a mutant delta-15 fatty acid desaturase. A plant in this embodiment may be soybean, oilseed *Brassica* species, sunflower, castor bean or corn. The mutant  
5 sequence may be derived from, for example, a *Brassica napus*, *Brassica rapa*, *Brassica juncea* or *Helianthus* delta-12 or delta-15 desaturase gene.

Another embodiment of the invention involves a method of producing a *Brassicaceae* or *Helianthus* plant line  
10 comprising the steps of: (a) inducing mutagenesis in cells of a starting variety of a *Brassicaceae* or *Helianthus* species; (b) obtaining progeny plants from the mutagenized cells; (c) identifying progeny plants that contain a mutation in a delta-12 or delta-15 fatty acid desaturase  
15 gene; and (d) producing a plant line by selfing or crossing. The resulting plant line may be subjected to mutagenesis in order to obtain a line having both a delta-12 desaturase mutation and a delta-15 desaturase mutation.

Yet another embodiment of the invention involves a  
20 method of producing plant lines containing altered fatty acid composition comprising: (a) crossing a first plant with a second plant having a mutant delta-12 or delta-15 fatty acid desaturase; (b) obtaining seeds from the cross of step (a); (c) growing fertile plants from such seeds; (d)  
25 obtaining progeny seed from the plants of step (c); and (e) identifying those seeds among the progeny that have altered fatty acid composition. Suitable plants are soybean, rapeseed, sunflower, safflower, castor bean and corn. Preferred plants are rapeseed and sunflower.

30 The invention is also embodied in vegetable oil obtained from plants disclosed herein, which vegetable oil has an altered fatty acid composition.

### Brief Description of the Sequence Listing

SEQ ID NO:1 shows a hypothetical DNA sequence of a *Brassica* Fad2 gene. SEQ ID NO:2 is the deduced amino acid sequence of SEQ ID NO:1.

5        SEQ ID NO:3 shows a hypothetical DNA sequence of a *Brassica* Fad2 gene having a mutation at nucleotide 316. SEQ ID NO:4 is the deduced amino acid sequence of SEQ ID NO:3.

10        SEQ ID NO:5 shows a hypothetical DNA sequence of a *Brassica* Fad2 gene. SEQ ID NO:6 is the deduced amino acid sequence of SEQ ID NO:5.

SEQ ID NO:7 shows a hypothetical DNA sequence of a *Brassica* Fad2 gene having a mutation at nucleotide 515. SEQ ID NO:8 is the deduced amino acid sequence of SEQ ID NO:7.

15        SEQ ID NO:9 shows the DNA sequence for the coding region of a wild type *Brassica* Fad2-D gene. SEQ ID NO:10 is the deduced amino acid sequence for SEQ ID NO:9.

SEQ ID NO:11 shows the DNA sequence for the coding region of the IMC 129 mutant *Brassica* Fad2-D gene. SEQ ID NO:12 is the deduced amino acid sequence for SEQ ID NO:11.

20        SEQ ID NO:13 shows the DNA sequence for the coding region of a wild type *Brassica* Fad2-F gene. SEQ ID NO:14 is the deduced amino acid sequence for SEQ ID NO:13.

25        SEQ ID NO:15 shows the DNA sequence for the coding region of the Q508 mutant *Brassica* Fad2-F gene. SEQ ID NO:16 is the deduced amino acid sequence for SEQ ID NO:15.

SEQ ID NO:17 shows the DNA sequence for the coding region of the Q4275 mutant *Brassica* Fad2-F gene. SEQ ID NO:18 is the deduced amino acid sequence for SEQ ID NO:17.

### Brief Description of the Figures

30        Figure 1 is a histogram showing the frequency distribution of seed oil oleic acid ( $C_{18:1}$ ) content in a segregating population of a Q508 X Westar cross. The bar

labeled WSGA 1A represents the  $C_{18:1}$  content of the Westar parent. The bar labeled Q508 represents the  $C_{18:1}$  content of the Q508 parent.

Sub  
C1  
5 Figure 2 shows the nucleotide sequences for a  
Brassica Fad2-D wild type gene (Fad2-D wt), IMC129 mutant  
gene (Fad2-D GA316 IMC129), Fad2-F wild type gene (Fad2-F  
wt), Q508 mutant gene (Fad2-F TA515 Q508) and Q4275 mutant  
gene (Fad2-F GA908 Q4275).

10 Figure 3 shows the deduced amino acid sequences for  
the polynucleotides of Figure 2.

#### Description of the Preferred Embodiments

All percent fatty acids herein are percent by weight  
of the oil of which the fatty acid is a component.

As used herein, a "line" is a group of plants that  
15 display little or no genetic variation between individuals  
for at least one trait. Such lines may be created by  
several generations of self-pollination and selection, or  
vegetative propagation from a single parent using tissue or  
cell culture techniques. As used herein, the term "variety"  
20 refers to a line which is used for commercial production.

The term "mutagenesis" refers to the use of a  
mutagenic agent to induce random genetic mutations within a  
population of individuals. The treated population, or a  
subsequent generation of that population, is then screened  
25 for usable trait(s) that result from the mutations. A  
"population" is any group of individuals that share a common  
gene pool. As used herein " $M_0$ " is untreated seed. As used  
herein, " $M_1$ " is the seed (and resulting plants) exposed to a  
mutagenic agent, while " $M_2$ " is the progeny (seeds and  
30 plants) of self-pollinated  $M_1$  plants, " $M_3$ " is the progeny of  
self-pollinated  $M_2$  plants, and " $M_4$ " is the progeny of self-  
pollinated  $M_3$  plants. " $M_5$ " is the progeny of self-

pollinated  $M_4$  plants. " $M_6$ ", " $M_7$ ", etc. are each the progeny of self-pollinated plants of the previous generation. The term "selfed" as used herein means self-pollinated.

"Stability" or "stable" as used herein means that  
5 with respect to a given fatty acid component, the component is maintained from generation to generation for at least two generations and preferably at least three generations at substantially the same level, e.g., preferably  $\pm 5\%$ . The method of invention is capable of creating lines with  
10 improved fatty acid compositions stable up to  $\pm 5\%$  from generation to generation. The above stability may be affected by temperature, location, stress and time of planting. Thus, comparison of fatty acid profiles should be made from seeds produced under similar growing conditions.  
15 Stability may be measured based on knowledge of prior generation.

Intensive breeding has produced *Brassica* plants whose seed oil contains less than 2% erucic acid. The same varieties have also been bred so that the defatted meal  
20 contains less than 30  $\mu\text{mol}$  glucosinolates/gram. "Canola" as used herein refers to plant variety seed or oil which contains less than 2% erucic acid ( $C_{22:1}$ ), and meal with less than 30  $\mu\text{mol}$  glucosinolates/gram.

Applicants have discovered plants with mutations in  
25 a delta-12 fatty acid desaturase gene. Such plants have useful alterations in the fatty acid compositions of the seed oil. Such mutations confer, for example, an elevated oleic acid content, a decreased, stabilized linoleic acid content, or both elevated oleic acid and decreased,  
30 stabilized linoleic acid content.

Applicants have further discovered plants with mutations in a delta-15 fatty acid desaturase gene. Such plants have useful alterations in the fatty acid composition



of the seed oil, e.g., a decreased, stabilized level of  $\alpha$ -linolenic acid.

Applicants have further discovered isolated nucleic acid fragments (polynucleotides) comprising sequences that carry mutations within the coding sequence of delta-12 or delta-15 fatty acid desaturases. The mutations confer desirable alterations in fatty acid levels in the seed oil of plants carrying such mutations. Delta-12 fatty acid desaturase is also known as omega-6 fatty acid desaturase and is sometimes referred to herein as Fad2 or 12-DES. Delta-15 fatty acid desaturase is also known <sup>as</sup> ~~on~~ omega-3 fatty acid desaturase and is sometimes referred to herein as Fad3 or 15-DES.

A nucleic acid fragment of the invention may be in the form of RNA or in the form of DNA, including cDNA, synthetic DNA or genomic DNA. The DNA may be double-stranded or single-stranded, and if single-stranded, can be either the coding strand or non-coding strand. An RNA analog may be, for example, mRNA or a combination of ribo- and deoxyribonucleotides. Illustrative examples of a nucleic acid fragment of the invention are the mutant sequences shown in Fig. 3.

A nucleic acid fragment of the invention contains a mutation in a microsomal delta-12 fatty acid desaturase coding sequence or a mutation in a microsomal delta-15 fatty acid desaturase coding sequence. Such a mutation renders the resulting desaturase gene product non-functional in plants, relative to the function of the gene product encoded by the wild-type sequence. The non-functionality of the delta-12 desaturase gene product can be inferred from the decreased level of reaction product (linoleic acid) and increased level of substrate (oleic acid) in plant tissues expressing the mutant sequence, compared to the

corresponding levels in plant tissues expressing the wild-type sequence. The non-functionality of the delta-15 desaturase gene product can be inferred from the decreased level of reaction product ( $\alpha$ -linolenic acid) and the increased level of substrate (linoleic acid) in plant tissues expressing the mutant sequence, compared to the corresponding levels in plant tissues expressing the wild-type sequence.

A nucleic acid fragment of the invention may comprise a portion of the coding sequence, e.g., at least about 10 nucleotides, provided that the fragment contains at least one mutation in the coding sequence. The length of a desired fragment depends upon the purpose for which the fragment will be used, e.g., PCR primer, site-directed mutagenesis and the like. In one embodiment, a nucleic acid fragment of the invention comprises the full length coding sequence of a mutant delta-12 or mutant delta-15 fatty acid desaturase, e.g., the mutant sequences of Fig. 3. In other embodiments, a nucleic acid fragment is about 20 to about 50 nucleotides (or base pairs, bp), or about 50 to about 500 nucleotides, or about 500 to about 1200 nucleotides in length.

A mutation in a nucleic acid fragment of the invention may be in any portion of the coding sequence that renders the resulting gene product non-functional. Suitable types of mutations include, without limitation, insertions of nucleotides, deletions of nucleotides, or transitions and transversions in the wild-type coding sequence. Such mutations result in insertions of one or more amino acids, deletions of one or more amino acids, and non-conservative amino acid substitutions in the corresponding gene product. In some embodiments, the sequence of a nucleic acid fragment

may comprise more than one mutation or more than one type of mutation.

Insertion or deletion of amino acids in a coding sequence may, for example, disrupt the conformation of essential alpha-helical or beta-pleated sheet regions of the resulting gene product. Amino acid insertions or deletions may also disrupt binding or catalytic sites important for gene product activity. It is known in the art that the insertion or deletion of a larger number of contiguous amino acids is more likely to render the gene product non-functional, compared to a smaller number of inserted or deleted amino acids.

Non-conservative amino acid substitutions may replace an amino acid of one class with an amino acid of a different class. Non-conservative substitutions may make a substantial change in the charge or hydrophobicity of the gene product. Non-conservative amino acid substitutions may also make a substantial change in the bulk of the residue side chain, e.g., substituting an alanyl residue for a isoleucyl residue.

Examples of non-conservative substitutions include the substitution of a basic amino acid for a non-polar amino acid, or a polar amino acid for an acidic amino acid. Because there are only 20 amino acids encoded in a gene, substitutions that result in a non-functional gene product may be determined by routine experimentation, incorporating amino acids of a different class in the region of the gene product targeted for mutation.

Preferred mutations are in a region of the nucleic acid encoding an amino acid sequence motif that is conserved among delta-12 fatty acid desaturases or delta-15 fatty acid desaturases, such as a His-Xaa-Xaa-Xaa-His motif (Tables 1-3). An example of a suitable region has a conserved HECGH

Sub  
C2

Sub  
C2  
correct

A

5 motif that is found, for example, in nucleotides  
corresponding to amino acids 105 to 109 of the *Arabidopsis*  
and *Brassica* delta-12 desaturase sequences, in nucleotides  
corresponding to amino acids 101 to 105 of the soybean  
delta-12 desaturase sequence and in nucleotides  
corresponding to amino acids 111 to 115 of the maize delta-  
12 desaturase sequence. See e.g., WO 94/11516<sup>11516</sup>; Okuley et  
al., *Plant Cell* 6:147-158 (1994). The one letter amino acid  
designations used herein are described in Alberts, B. et  
10 al., *Molecular Biology of the Cell*, 3rd edition, Garland  
Publishing, New York, 1994. Amino acids flanking this motif  
are also highly conserved among delta-12 and delta-15  
desaturases and are also suitable candidates for mutations  
in fragments of the invention.

15 An illustrative embodiment of a mutation in a  
nucleic acid fragment of the invention is a Glu to Lys  
substitution in the HECGH motif of a *Brassica* microsomal  
delta-12 desaturase sequence, either the D form or the F  
form. This mutation results in the sequence HECGH being  
20 changed to HECGH as seen by comparing SEQ ID NO:10 (wild-  
type D form) to SEQ ID NO:12 (mutant D form). A similar  
mutation in other Fad-2 sequences is contemplated to result  
in a non-functional gene product. (Compare SEQ ID NO:2 to  
SEQ ID NO:4)

25 A similar motif may be found at amino acids 101 to  
105 of the *Arabidopsis* microsomal delta-15 fatty acid  
desaturase, as well as in the corresponding rape and soybean  
desaturases (Table 5). See, e.g., WO 93/11245; Arondel, V.  
et al., *Science*, 258:1153-1155 (1992); Yadav, N. et al.,  
30 *Plant Physiol.*, 103:467-476 (1993). Plastid delta-15 fatty  
acids have a similar motif (Table 5).

Sub  
C3

Among the types of mutations in an HECGH motif that  
render the resulting gene product non-functional are non-

Sub  
C3  
C4  
C5  
C6  
C7  
C8  
C9  
C10  
C11  
C12  
C13  
C14  
C15  
C16  
C17  
C18  
C19  
C20  
C21  
C22  
C23  
C24  
C25  
C26  
C27  
C28  
C29  
C30  
C31  
C32  
C33  
C34  
C35  
C36  
C37  
C38  
C39  
C40  
C41  
C42  
C43  
C44  
C45  
C46  
C47  
C48  
C49  
C50  
C51  
C52  
C53  
C54  
C55  
C56  
C57  
C58  
C59  
C60  
C61  
C62  
C63  
C64  
C65  
C66  
C67  
C68  
C69  
C70  
C71  
C72  
C73  
C74  
C75  
C76  
C77  
C78  
C79  
C80  
C81  
C82  
C83  
C84  
C85  
C86  
C87  
C88  
C89  
C90  
C91  
C92  
C93  
C94  
C95  
C96  
C97  
C98  
C99  
C100  
C101  
C102  
C103  
C104  
C105  
C106  
C107  
C108  
C109  
C110  
C111  
C112  
C113  
C114  
C115  
C116  
C117  
C118  
C119  
C120  
C121  
C122  
C123  
C124  
C125  
C126  
C127  
C128  
C129  
C130  
C131  
C132  
C133  
C134  
C135  
C136  
C137  
C138  
C139  
C140  
C141  
C142  
C143  
C144  
C145  
C146  
C147  
C148  
C149  
C150  
C151  
C152  
C153  
C154  
C155  
C156  
C157  
C158  
C159  
C160  
C161  
C162  
C163  
C164  
C165  
C166  
C167  
C168  
C169  
C170  
C171  
C172  
C173  
C174  
C175  
C176  
C177  
C178  
C179  
C180  
C181  
C182  
C183  
C184  
C185  
C186  
C187  
C188  
C189  
C190  
C191  
C192  
C193  
C194  
C195  
C196  
C197  
C198  
C199  
C200  
C201  
C202  
C203  
C204  
C205  
C206  
C207  
C208  
C209  
C210  
C211  
C212  
C213  
C214  
C215  
C216  
C217  
C218  
C219  
C220  
C221  
C222  
C223  
C224  
C225  
C226  
C227  
C228  
C229  
C230  
C231  
C232  
C233  
C234  
C235  
C236  
C237  
C238  
C239  
C240  
C241  
C242  
C243  
C244  
C245  
C246  
C247  
C248  
C249  
C250  
C251  
C252  
C253  
C254  
C255  
C256  
C257  
C258  
C259  
C260  
C261  
C262  
C263  
C264  
C265  
C266  
C267  
C268  
C269  
C270  
C271  
C272  
C273  
C274  
C275  
C276  
C277  
C278  
C279  
C280  
C281  
C282  
C283  
C284  
C285  
C286  
C287  
C288  
C289  
C290  
C291  
C292  
C293  
C294  
C295  
C296  
C297  
C298  
C299  
C300  
C301  
C302  
C303  
C304  
C305  
C306  
C307  
C308  
C309  
C310  
C311  
C312  
C313  
C314  
C315  
C316  
C317  
C318  
C319  
C320  
C321  
C322  
C323  
C324  
C325  
C326  
C327  
C328  
C329  
C330  
C331  
C332  
C333  
C334  
C335  
C336  
C337  
C338  
C339  
C340  
C341  
C342  
C343  
C344  
C345  
C346  
C347  
C348  
C349  
C350  
C351  
C352  
C353  
C354  
C355  
C356  
C357  
C358  
C359  
C360  
C361  
C362  
C363  
C364  
C365  
C366  
C367  
C368  
C369  
C370  
C371  
C372  
C373  
C374  
C375  
C376  
C377  
C378  
C379  
C380  
C381  
C382  
C383  
C384  
C385  
C386  
C387  
C388  
C389  
C390  
C391  
C392  
C393  
C394  
C395  
C396  
C397  
C398  
C399  
C400  
C401  
C402  
C403  
C404  
C405  
C406  
C407  
C408  
C409  
C410  
C411  
C412  
C413  
C414  
C415  
C416  
C417  
C418  
C419  
C420  
C421  
C422  
C423  
C424  
C425  
C426  
C427  
C428  
C429  
C430  
C431  
C432  
C433  
C434  
C435  
C436  
C437  
C438  
C439  
C440  
C441  
C442  
C443  
C444  
C445  
C446  
C447  
C448  
C449  
C450  
C451  
C452  
C453  
C454  
C455  
C456  
C457  
C458  
C459  
C460  
C461  
C462  
C463  
C464  
C465  
C466  
C467  
C468  
C469  
C470  
C471  
C472  
C473  
C474  
C475  
C476  
C477  
C478  
C479  
C480  
C481  
C482  
C483  
C484  
C485  
C486  
C487  
C488  
C489  
C490  
C491  
C492  
C493  
C494  
C495  
C496  
C497  
C498  
C499  
C500  
C501  
C502  
C503  
C504  
C505  
C506  
C507  
C508  
C509  
C510  
C511  
C512  
C513  
C514  
C515  
C516  
C517  
C518  
C519  
C520  
C521  
C522  
C523  
C524  
C525  
C526  
C527  
C528  
C529  
C530  
C531  
C532  
C533  
C534  
C535  
C536  
C537  
C538  
C539  
C540  
C541  
C542  
C543  
C544  
C545  
C546  
C547  
C548  
C549  
C550  
C551  
C552  
C553  
C554  
C555  
C556  
C557  
C558  
C559  
C560  
C561  
C562  
C563  
C564  
C565  
C566  
C567  
C568  
C569  
C570  
C571  
C572  
C573  
C574  
C575  
C576  
C577  
C578  
C579  
C580  
C581  
C582  
C583  
C584  
C585  
C586  
C587  
C588  
C589  
C590  
C591  
C592  
C593  
C594  
C595  
C596  
C597  
C598  
C599  
C600  
C601  
C602  
C603  
C604  
C605  
C606  
C607  
C608  
C609  
C610  
C611  
C612  
C613  
C614  
C615  
C616  
C617  
C618  
C619  
C620  
C621  
C622  
C623  
C624  
C625  
C626  
C627  
C628  
C629  
C630  
C631  
C632  
C633  
C634  
C635  
C636  
C637  
C638  
C639  
C640  
C641  
C642  
C643  
C644  
C645  
C646  
C647  
C648  
C649  
C650  
C651  
C652  
C653  
C654  
C655  
C656  
C657  
C658  
C659  
C660  
C661  
C662  
C663  
C664  
C665  
C666  
C667  
C668  
C669  
C670  
C671  
C672  
C673  
C674  
C675  
C676  
C677  
C678  
C679  
C680  
C681  
C682  
C683  
C684  
C685  
C686  
C687  
C688  
C689  
C690  
C691  
C692  
C693  
C694  
C695  
C696  
C697  
C698  
C699  
C700  
C701  
C702  
C703  
C704  
C705  
C706  
C707  
C708  
C709  
C710  
C711  
C712  
C713  
C714  
C715  
C716  
C717  
C718  
C719  
C720  
C721  
C722  
C723  
C724  
C725  
C726  
C727  
C728  
C729  
C730  
C731  
C732  
C733  
C734  
C735  
C736  
C737  
C738  
C739  
C740  
C741  
C742  
C743  
C744  
C745  
C746  
C747  
C748  
C749  
C750  
C751  
C752  
C753  
C754  
C755  
C756  
C757  
C758  
C759  
C760  
C761  
C762  
C763  
C764  
C765  
C766  
C767  
C768  
C769  
C770  
C771  
C772  
C773  
C774  
C775  
C776  
C777  
C778  
C779  
C780  
C781  
C782  
C783  
C784  
C785  
C786  
C787  
C788  
C789  
C790  
C791  
C792  
C793  
C794  
C795  
C796  
C797  
C798  
C799  
C800  
C801  
C802  
C803  
C804  
C805  
C806  
C807  
C808  
C809  
C810  
C811  
C812  
C813  
C814  
C815  
C816  
C817  
C818  
C819  
C820  
C821  
C822  
C823  
C824  
C825  
C826  
C827  
C828  
C829  
C830  
C831  
C832  
C833  
C834  
C835  
C836  
C837  
C838  
C839  
C840  
C841  
C842  
C843  
C844  
C845  
C846  
C847  
C848  
C849  
C850  
C851  
C852  
C853  
C854  
C855  
C856  
C857  
C858  
C859  
C860  
C861  
C862  
C863  
C864  
C865  
C866  
C867  
C868  
C869  
C870  
C871  
C872  
C873  
C874  
C875  
C876  
C877  
C878  
C879  
C880  
C881  
C882  
C883  
C884  
C885  
C886  
C887  
C888  
C889  
C890  
C891  
C892  
C893  
C894  
C895  
C896  
C897  
C898  
C899  
C900  
C901  
C902  
C903  
C904  
C905  
C906  
C907  
C908  
C909  
C910  
C911  
C912  
C913  
C914  
C915  
C916  
C917  
C918  
C919  
C920  
C921  
C922  
C923  
C924  
C925  
C926  
C927  
C928  
C929  
C930  
C931  
C932  
C933  
C934  
C935  
C936  
C937  
C938  
C939  
C940  
C941  
C942  
C943  
C944  
C945  
C946  
C947  
C948  
C949  
C950  
C951  
C952  
C953  
C954  
C955  
C956  
C957  
C958  
C959  
C960  
C961  
C962  
C963  
C964  
C965  
C966  
C967  
C968  
C969  
C970  
C971  
C972  
C973  
C974  
C975  
C976  
C977  
C978  
C979  
C980  
C981  
C982  
C983  
C984  
C985  
C986  
C987  
C988  
C989  
C990  
C991  
C992  
C993  
C994  
C995  
C996  
C997  
C998  
C999  
C1000  
C1001  
C1002  
C1003  
C1004  
C1005  
C1006  
C1007  
C1008  
C1009  
C1010  
C1011  
C1012  
C1013  
C1014  
C1015  
C1016  
C1017  
C1018  
C1019  
C1020  
C1021  
C1022  
C1023  
C1024  
C1025  
C1026  
C1027  
C1028  
C1029  
C1030  
C1031  
C1032  
C1033  
C1034  
C1035  
C1036  
C1037  
C1038  
C1039  
C1040  
C1041  
C1042  
C1043  
C1044  
C1045  
C1046  
C1047  
C1048  
C1049  
C1050  
C1051  
C1052  
C1053  
C1054  
C1055  
C1056  
C1057  
C1058  
C1059  
C1060  
C1061  
C1062  
C1063  
C1064  
C1065  
C1066  
C1067  
C1068  
C1069  
C1070  
C1071  
C1072  
C1073  
C1074  
C1075  
C1076  
C1077  
C1078  
C1079  
C1080  
C1081  
C1082  
C1083  
C1084  
C1085  
C1086  
C1087  
C1088  
C1089  
C1090  
C1091  
C1092  
C1093  
C1094  
C1095  
C1096  
C1097  
C1098  
C1099  
C1100  
C1101  
C1102  
C1103  
C1104  
C1105  
C1106  
C1107  
C1108  
C1109  
C1110  
C1111  
C1112  
C1113  
C1114  
C1115  
C1116  
C1117  
C1118  
C1119  
C1120  
C1121  
C1122  
C1123  
C1124  
C1125  
C1126  
C1127  
C1128  
C1129  
C1130  
C1131  
C1132  
C1133  
C1134  
C1135  
C1136  
C1137  
C1138  
C1139  
C1140  
C1141  
C1142  
C1143  
C1144  
C1145  
C1146  
C1147  
C1148  
C1149  
C1150  
C1151  
C1152  
C1153  
C1154  
C1155  
C1156  
C1157  
C1158  
C1159  
C1160  
C1161  
C1162  
C1163  
C1164  
C1165  
C1166  
C1167  
C1168  
C1169  
C1170  
C1171  
C1172  
C1173  
C1174  
C1175  
C1176  
C1177  
C1178  
C1179  
C1180  
C1181  
C1182  
C1183  
C1184  
C1185  
C1186  
C1187  
C1188  
C1189  
C1190  
C1191  
C1192  
C1193  
C1194  
C1195  
C1196  
C1197  
C1198  
C1199  
C1200  
C1201  
C1202  
C1203  
C1204  
C1205  
C1206  
C1207  
C1208  
C1209  
C1210  
C1211  
C1212  
C1213  
C1214  
C1215  
C1216  
C1217  
C1218  
C1219  
C1220  
C1221  
C1222  
C1223  
C1224  
C1225  
C1226  
C1227  
C1228  
C1229  
C1230  
C1231  
C1232  
C1233  
C1234  
C1235  
C1236  
C1237  
C1238  
C1239  
C1240  
C1241  
C1242  
C1243  
C1244  
C1245  
C1246  
C1247  
C1248  
C1249  
C1250  
C1251  
C1252  
C1253  
C1254  
C1255  
C1256  
C1257  
C1258  
C1259  
C1260  
C1261  
C1262  
C1263  
C1264  
C1265  
C1266  
C1267  
C1268  
C1269  
C1270  
C1271  
C1272  
C1273  
C1274  
C1275  
C1276  
C1277  
C1278  
C1279  
C1280  
C1281  
C1282  
C1283  
C1284  
C1285  
C1286  
C1287  
C1288  
C1289  
C1290  
C1291  
C1292  
C1293  
C1294  
C1295  
C1296  
C1297  
C1298  
C1299  
C1300  
C1301  
C1302  
C1303  
C1304  
C1305  
C1306  
C1307  
C1308  
C1309  
C1310  
C1311  
C1312  
C1313  
C1314  
C1315  
C1316  
C1317  
C1318  
C1319  
C1320  
C1321  
C1322  
C1323  
C1324  
C1325  
C1326  
C1327  
C1328  
C1329  
C1330  
C1331  
C1332  
C1333  
C1334  
C1335  
C1336  
C1337  
C1338  
C1339  
C1340  
C1341  
C1342  
C1343  
C1344  
C1345  
C1346  
C1347  
C1348  
C1349  
C1350  
C1351  
C1352  
C1353  
C1354  
C1355  
C1356  
C1357  
C1358  
C1359  
C1360  
C1361  
C1362  
C1363  
C1364  
C1365  
C1366  
C1367  
C1368  
C1369  
C1370  
C1371  
C1372  
C1373  
C1374  
C1375  
C1376  
C1377  
C1378  
C1379  
C1380  
C1381  
C1382  
C1383  
C1384  
C1385  
C1386  
C1387  
C1388  
C1389  
C1390  
C1391  
C1392  
C1393  
C1394  
C1395  
C1396  
C1397  
C1398  
C1399  
C1400  
C1401  
C1402  
C1403  
C1404  
C1405  
C1406  
C1407  
C1408  
C1409  
C1410  
C1411  
C1412  
C1413  
C1414  
C1415  
C1416  
C1417  
C1418  
C1419  
C1420  
C1421  
C1422  
C1423  
C1424  
C1425  
C1426  
C1427  
C1428  
C1429  
C1430  
C1431  
C1432  
C1433  
C1434  
C1435  
C1436  
C1437  
C1438  
C1439  
C1440  
C1441  
C1442  
C1443  
C1444  
C1445  
C1446  
C1447  
C1448  
C1449  
C1450  
C1451  
C1452  
C1453  
C1454  
C1455  
C1456  
C1457  
C1458  
C1459  
C1460  
C1461  
C1462  
C1463  
C1464  
C1465  
C1466  
C1467  
C1468  
C1469  
C1470  
C1471  
C1472  
C1473  
C1474  
C1475  
C1476  
C1477  
C1478  
C1479  
C1480  
C1481  
C1482  
C1483  
C1484  
C1485  
C1486  
C1487  
C1488  
C1489  
C1490  
C1491  
C1492  
C1493  
C1494  
C1495  
C1496  
C1497  
C1498  
C1499  
C1500  
C1501  
C1502  
C1503  
C1504  
C1505  
C1506  
C1507  
C1508  
C1509  
C1510  
C1511  
C1512  
C1513  
C1514  
C1515  
C1516  
C1517  
C1518  
C1519  
C1520  
C1521  
C1522  
C1523  
C1524  
C1525  
C1526  
C1527  
C1528  
C1529  
C1530  
C1531  
C1532  
C1533  
C1534  
C1535  
C1536  
C1537  
C1538  
C1539  
C1540  
C1541  
C1542  
C1543  
C1544  
C1545  
C1546  
C1547  
C1548  
C1549  
C1550  
C1551  
C1552  
C1553  
C1554  
C1555  
C1556  
C1557  
C1558  
C1559  
C1560  
C1561  
C1562  
C1563  
C1564  
C1565  
C1566  
C1567  
C1568  
C1569  
C1570  
C1571  
C1572  
C1573  
C1574  
C1575  
C1576  
C1577  
C1578  
C1579  
C1580  
C1581  
C1582  
C1583  
C1584  
C1585  
C1586  
C1587  
C1588  
C1589  
C1590  
C1591  
C1592  
C1593  
C1594  
C1595  
C1596  
C1597  
C1598  
C1599  
C1600  
C1601  
C1602  
C1603  
C1604  
C1605  
C1606  
C1607  
C1608  
C1609  
C1610  
C1611  
C1612  
C1613  
C1614  
C1615  
C1616  
C1617  
C1618  
C1619  
C1620  
C1621  
C1622  
C1623  
C1624  
C1625  
C1626  
C1627  
C1628  
C1629  
C1630  
C1631  
C1632  
C1633  
C1634  
C1635  
C1636  
C1637  
C1638  
C1639  
C1640  
C1641  
C1642  
C1643  
C1644  
C1645  
C1646  
C1647  
C1648  
C1649  
C1650  
C1651  
C1652  
C1653  
C1654  
C1655  
C1656  
C1657  
C1658  
C1659  
C1660  
C1661  
C1662  
C1663  
C1664  
C1665  
C1666  
C1667  
C1668  
C1669  
C1670  
C1671  
C1672  
C1673  
C1674  
C1675  
C1676  
C1677  
C1678  
C1679  
C1680  
C1681  
C1682  
C1683  
C1684  
C1685  
C1686  
C1687  
C1688  
C1689  
C1690  
C1691  
C1692  
C1693  
C1694  
C1695  
C1696  
C1697  
C1698  
C1699  
C1700  
C1701  
C1702  
C1703  
C1704  
C1705  
C1706  
C1707  
C1708  
C1709  
C1710  
C1711  
C1712  
C1713  
C1714  
C1715  
C1716  
C1717  
C1718  
C1719  
C1720  
C1721  
C1722  
C1723  
C1724  
C1725  
C1726  
C1727  
C1728  
C1729  
C1730  
C1731  
C1732  
C1733  
C1734  
C1735  
C1736  
C1737  
C1738  
C1739  
C1740  
C1741  
C1742  
C1743  
C1744  
C1745  
C1746  
C1747  
C1748  
C1749  
C1750  
C1751  
C1752  
C1753  
C1754  
C1755  
C1756  
C1757  
C1758  
C1759  
C1760  
C1761  
C1762  
C1763  
C1764  
C1765  
C1766  
C1767  
C1768  
C1769  
C1770  
C1771  
C1772  
C1773  
C1774  
C1775  
C1776  
C1777  
C1778  
C1779  
C1780  
C1781  
C1782  
C1783  
C1784  
C1785  
C1786  
C1787  
C1788  
C1789  
C1790  
C1791  
C1792  
C1793  
C1794  
C1795  
C1796  
C1797  
C1798  
C1799  
C1800  
C1801  
C1802  
C1803  
C1804  
C1805  
C1806  
C1807  
C1808  
C1809  
C1810  
C1811  
C1812  
C1813  
C1814  
C1815  
C1816  
C1817  
C1818  
C1819  
C1820  
C1821  
C1822  
C1823  
C1824  
C1825  
C1826  
C1827  
C1828  
C1829  
C1830  
C1831  
C1832  
C1833  
C1834  
C1835  
C1836  
C1837  
C1838  
C1839  
C1840  
C1841  
C1842  
C1843  
C1844  
C1845  
C1846  
C1847  
C1848  
C1849  
C1850  
C1851  
C1852  
C1853  
C1854  
C1855  
C1856  
C1857  
C1858  
C1859  
C1860  
C1861  
C1862  
C1863  
C1864  
C1865  
C1866  
C1867  
C1868  
C1869  
C1870  
C1871  
C1872  
C1873  
C1874  
C1875  
C1876  
C1877  
C1878  
C1879  
C1880  
C1881  
C1882  
C1883  
C1884  
C1885  
C1886  
C1887  
C1888  
C1889  
C1890  
C1891  
C1892  
C1893  
C1894  
C1895  
C1896  
C1897  
C1898  
C1899  
C1900  
C1901  
C1902  
C1903  
C1904  
C1905  
C1906  
C1907  
C1908  
C1909  
C1910  
C1911  
C1912  
C1913  
C1914  
C1915  
C1916  
C1917  
C1918  
C1919  
C1920  
C1921  
C1922  
C1923  
C1924

TABLE 1

Alignment of Amino Acid Sequences from Microsomal  
Delta-12 Fatty Acid Desaturases

Sub  
CS >

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>Arabidopsis thaliana</i>	100-129	IWVIAHECGH HAFSDYQWLD DTVGLIFHSF
	<i>Glycine max</i>	96-125	VWVIAHECGH HAFSKYQWVD DVVGLTLHST
	<i>Zea mays</i>	106-135	VWVIAHECGH HAFSDYSLLD DVVGLVLHSS
	<i>Ricinus communis</i> <sup>a</sup>	1- 29	WVMAHDCGH HAFSDYQLLD DVVGLILHSC
	<i>Brassica napus D</i>	100-128	VWVIAHECGH HAFSDYQWLD DTVGLIFHS
10	<i>Brassica napus F</i>	100-128	VWVIAHECGH HAFSDYQWLD DTVGLIFHS

<sup>a</sup> from plasmid pRF2-1C

TABLE 2

Alignment of Amino Acid Sequences from Microsomal  
Delta-12 Fatty Acid Desaturases

09771904-012901

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
15	<i>Arabidopsis thaliana</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Glycine max</i>	126-154	LLVPYFSWKI SHRRHHSNTG SLDRDEVFV
	<i>Zea mays</i>	136-164	LMVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Ricinus communis</i> <sup>a</sup>	30- 58	LLVPYFSWKH SHRRHHSNTG SLERDEVFV
20	<i>Brassica napus D</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Brassica napus F</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV

<sup>a</sup> from plasmid pRF2-1C

TABLE 3

Alignment of Amino Acid Sequences from Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
25	<i>Arabidopsis thaliana</i>	298-333	DRDYGILNKV FHNITDTHVA HHLFSTMPHY NAMEAT
	<i>Glycine max</i>	294-329	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT
	<i>Zea mays</i>	305-340	DRDYGILNRV FHNITDTHVA HHLFSTMPHY HAMEAT
30	<i>Ricinus communis</i> <sup>a</sup>	198-224	DRDYGILNKV FHNITDTQVA HHLF TMP
	<i>Brassica napus D</i>	299-334	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT
	<i>Brassica napus F</i>	299-334	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT

<sup>a</sup> from plasmid pRF2-1C

TABLE 4

Alignment of Conserved Amino Acids from Microsomal  
Delta-12 Fatty Acid Desaturases

Sub  
C5  
amold

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>Arabidopsis thaliana</i>	165-180	IKWYGKYLNN PLGRIM
	<i>Glycine max</i>	161-176	VAWFSLYLNN PLGRAV
	<i>Zea mays</i>	172-187	PWYTPYVYNN PVGRVV
	<i>Ricinus communis</i> <sup>a</sup>	65-80	IRWYSKYLNN PPGRIM
	<i>Brassica napus</i> D	165-180	IKWYGKYLNN PLGRTV
10	<i>Brassica napus</i> F	165-180	IKWYGKYLNN PLGRTV

<sup>a</sup> from plasmid pRF2-1C

TABLE 5

Alignment of Conserved Amino Acids from Plastid and Microsomal  
Delta-15 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
15	<i>Arabidopsis thaliana</i> <sup>a</sup>	156-177	WALFVLGHD CGHGSFSNDP KLN
	<i>Brassica napus</i> <sup>a</sup>	114-135	WALFVLGHD CGHGSFSNDP RLN
	<i>Glycine max</i> <sup>a</sup>	164-185	WALFVLGHD CGHGSFSNNS KLN
	<i>Arabidopsis thaliana</i>	94-115	WAIFVLGHD CGHGSFSDIP LLN
20	<i>Brassica napus</i>	87-109	WALFVLGHD CGHGSFSNDP RLN
	<i>Glycine max</i>	93-114	WALFVLGHD CGHGSFSDSP PLN

<sup>a</sup> Plastid sequences

TABLE 6

Alignment of Conserved Amino Acids from Plastid and Microsomal  
Delta-15 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
25	<i>A. thaliana</i> <sup>a</sup>	188-216	ILVPYHGWRI SHRTHHQNHG HVENDESWH
	<i>B. napus</i> <sup>a</sup>	146-174	ILVPYHGWRI SHRTHHQNHG HVENDESWH
	<i>Glycine max</i> <sup>a</sup>	196-224	ILVPYHGWRI SHRTHHQNHG HAENDESWH
30	<i>A. thaliana</i>	126-154	ILVPYHGWRI SHRTHHQNHG HVENDESWV
	<i>Brassica napus</i>	117-145	ILVPYHGWRI SHRTHHQNHG HVENDESWV
	<i>Glycine max</i>	125-153	ILVPYHGWRI SHRTHHQNHG HIEKDESWV

<sup>a</sup> Plastid sequences

The conservation of amino acid motifs and their  
relative positions indicates that regions of a delta-12 or

delta-15 fatty acid desaturase that can be mutated in one species to generate a non-functional desaturase can be mutated in the corresponding region from other species to generate a non-functional delta-12 desaturase or delta-15 desaturase gene product in that species.

Mutations in any of the regions of Tables 1-6 are specifically included within the scope of the invention and are substantially identical to those mutations exemplified herein, provided that such mutation (or mutations) renders the resulting desaturase gene product non-functional, as discussed hereinabove.

A nucleic acid fragment containing a mutant sequence can be generated by techniques known to the skilled artisan. Such techniques include, without limitation, site-directed mutagenesis of wild-type sequences and direct synthesis using automated DNA synthesizers.

A nucleic acid fragment containing a mutant sequence can also be generated by mutagenesis of plant seeds or regenerable plant tissue by, e.g., ethyl methane sulfonate, X-rays or other mutagens. With mutagenesis, mutant plants having the desired fatty acid phenotype in seeds are identified by known techniques and a nucleic acid fragment containing the desired mutation is isolated from genomic DNA or RNA of the mutant line. The site of the specific mutation is then determined by sequencing the coding region of the delta-12 desaturase or delta-15 desaturase gene. Alternatively, labeled nucleic acid probes that are specific for desired mutational events can be used to rapidly screen a mutagenized population.

The disclosed method may be applied to all oilseed Brassica species, and to both Spring and Winter maturing types within each species. Physical mutagens, including but not limited to X-rays, UV rays, and other physical



0974904-012901  
treatments which cause chromosome damage, and other chemical mutagens, including but not limited to ethidium bromide, nitrosoguanidine, diepoxybutane etc. may also be used to induce mutations. The mutagenesis treatment may also be applied to other stages of plant development, including but not limited to cell cultures, embryos, microspores and shoot apices.

"Stable mutations" as used herein are defined as  $M_s$  or more advanced lines which maintain a selected altered fatty acid profile for a minimum of three generations, including a minimum of two generations under field conditions, and exceeding established statistical thresholds for a minimum of two generations, as determined by gas chromatographic analysis of a minimum of 10 randomly selected seeds bulked together. Alternatively, stability may be measured in the same way by comparing to subsequent generations. In subsequent generations, stability is defined as having similar fatty acid profiles in the seed as that of the prior or subsequent generation when grown under substantially similar conditions.

Mutation breeding has traditionally produced plants carrying, in addition to the trait of interest, multiple, deleterious traits, e.g., reduced plant vigor and reduced fertility. Such traits may indirectly affect fatty acid composition, producing an unstable mutation; and/or reduce yield, thereby reducing the commercial utility of the invention. To eliminate the occurrence of deleterious mutations and reduce the load of mutations carried by the plant, a low mutagen dose is used in the seed treatments to create an LD30 population. This allows for the rapid selection of single gene mutations for fatty acid traits in agronomic backgrounds which produce acceptable yields.

Sub  
C6 >  
The seeds of several different fatty acid lines have been deposited with the American Type Culture Collection and have the following accession numbers.

	<u>Line</u>	<u>Accession No.</u>	<u>Deposit Date</u>
5	A129.5	40811	May 25, 1990
	A133.1	40812	May 25, 1990
	M3032.1	75021	June 7, 1991
	M3062.8	75025	June 7, 1991
	M3028.10	75026	June 7, 1991
10	IMC130	75446	April 16, 1993
	Q4275	97569	May 10, 1996

09771904-012901  
In some plant species or varieties more than one form of endogenous microsomal delta-12 desaturase may be found. In amphidiploids, each form may be derived from one of the parent genomes making up the species under consideration. Plants with mutations in both forms have a fatty acid profile that differs from plants with a mutation in only one form. An example of such a plant is *Brassica napus* line Q508, a doubly-mutagenized line containing a mutant D-form of delta-12 desaturase (SEQ ID NO:11) and a mutant F-form of delta-12 desaturase (SEQ ID NO:15). Another example is line Q4275, which contains a mutant D-form of delta-12 desaturase (SEQ ID NO:11) and a mutant F-form of delta-12 desaturase (SEQ ID NO:17). See Figs. 2-3.

Preferred host or recipient organisms for introduction of a nucleic acid fragment of the invention are the oil-producing species, such as soybean (*Glycine max*), rapeseed (e.g., *Brassica napus*, *B. rapa* and *B. juncea*), sunflower (*Helianthus annuus*), castor bean (*Ricinus communis*), corn (*Zea mays*), and safflower (*Carthamus tinctorius*).

A nucleic acid fragment of the invention may further comprise additional nucleic acids. For example, a nucleic acid encoding a secretory or leader amino acid sequence can

be linked to a mutant desaturase nucleic acid fragment such that the secretory or leader sequence is fused in-frame to the amino terminal end of a mutant delta-12 or delta-15 desaturase polypeptide. Other nucleic acid fragments are known in the art that encode amino acid sequences useful for fusing in-frame to the mutant desaturase polypeptides disclosed herein. See, e.g., U.S. 5,629,193 incorporated herein by reference. A nucleic acid fragment may also have one or more regulatory elements operably linked thereto.

The present invention also comprises nucleic acid fragments that selectively hybridize to mutant desaturase sequences. Such a nucleic acid fragment typically is at least 15 nucleotides in length. Hybridization typically involves Southern analysis (Southern blotting), a method by which the presence of DNA sequences in a target nucleic acid mixture are identified by hybridization to a labeled oligonucleotide or DNA fragment probe. Southern analysis typically involves electrophoretic separation of DNA digests on agarose gels, denaturation of the DNA after electrophoretic separation, and transfer of the DNA to nitrocellulose, nylon, or another suitable membrane support for analysis with a radiolabeled, biotinylated, or enzyme-labeled probe as described in sections 9.37-9.52 of Sambrook et al., (1989) *Molecular Cloning*, second edition, Cold Spring Harbor Laboratory, Plainview; NY.

A nucleic acid fragment can hybridize under moderate stringency conditions or, preferably, under high stringency conditions to a mutant desaturase sequence. High stringency conditions are used to identify nucleic acids that have a high degree of homology to the probe. High stringency conditions can include the use of low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate (0.1X SSC); 0.1% sodium lauryl

09771904-012901

sulfate (SDS) at 50-65°C. Alternatively, a denaturing agent such as formamide can be employed during hybridization, e.g., 50% formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42°C. Another example is the use of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS.

Moderate stringency conditions refers to hybridization conditions used to identify nucleic acids that have a lower degree of identity to the probe than do nucleic acids identified under high stringency conditions. Moderate stringency conditions can include the use of higher ionic strength and/or lower temperatures for washing of the hybridization membrane, compared to the ionic strength and temperatures used for high stringency hybridization. For example, a wash solution comprising 0.060 M NaCl/0.0060 M sodium citrate (4X SSC) and 0.1% sodium lauryl sulfate (SDS) can be used at 50°C, with a last wash in 1X SSC, at 65°C. Alternatively, a hybridization wash in 1X SSC at 37°C can be used.

Hybridization can also be done by Northern analysis (Northern blotting), a method used to identify RNAs that hybridize to a known probe such as an oligonucleotide, DNA fragment, cDNA or fragment thereof, or RNA fragment. The probe is labeled with a radioisotope such as <sup>32</sup>P, by biotinylation or with an enzyme. The RNA to be analyzed can be usually electrophoretically separated on an agarose or polyacrylamide gel, transferred to nitrocellulose, nylon, or other suitable membrane, and hybridized with the probe,

using standard techniques well known in the art such as those described in sections 7.39-7.52 of Sambrook et al., *supra*.

5 A polypeptide of the invention comprises an isolated polypeptide having a mutant amino acid sequence, as well as derivatives and analogs thereof. See, e.g., the mutant amino acid sequences of Fig. 3. By "isolated" is meant a polypeptide that is expressed and produced in an environment other than the environment in which the polypeptide is  
10 naturally expressed and produced. For example, a plant polypeptide is isolated when expressed and produced in bacteria or fungi. A polypeptide of the invention also comprises variants of the mutant desaturase polypeptides disclosed herein, as discussed above.

15 In one embodiment of the claimed invention, a plant contains both a delta-12 desaturase mutation and a delta-15 desaturase mutation. Such plants can have a fatty acid composition comprising very high oleic acid and very low alpha-linolenic acid levels. Mutations in delta-12  
20 desaturase and delta-15 desaturase may be combined in a plant by making a genetic cross between delta-12 desaturase and delta-15 desaturase single mutant lines. A plant having a mutation in delta-12 fatty acid desaturase is crossed or mated with a second plant having a mutation in delta-15  
25 fatty acid desaturase. Seeds produced from the cross are planted and the resulting plants are selfed in order to obtain progeny seeds. These progeny seeds are then screened in order to identify those seeds carrying both mutant genes.

Alternatively, a line possessing either a delta-12  
30 desaturase or a delta-15 desaturase mutation can be subjected to mutagenesis to generate a plant or plant line having mutations in both delta-12 desaturase and delta-15 desaturase. For example, the IMC 129 line has a mutation in

the coding region (Glu<sub>106</sub> to Lys<sub>106</sub>) of the D form of the  
microsomal delta-12 desaturase structural gene. Cells  
(e.g., seeds) of this line can be mutagenized to induce a  
mutation in a delta-15 desaturase gene, resulting in a plant  
5 or plant line carrying a mutation in a delta-12 fatty acid  
desaturase gene and a mutation in a delta-15 fatty acid  
desaturase gene.

Progeny includes descendants of a particular plant  
or plant line, e.g., seeds developed on an instant plant are  
10 descendants. Progeny of an instant plant include seeds  
formed on F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, and subsequent generation plants, or  
seeds formed on BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub>, and subsequent generation  
plants.

Plants according to the invention preferably contain  
15 an altered fatty acid composition. For example, oil  
obtained from seeds of such plants may have from about 69 to  
about 90% oleic acid, based on the total fatty acid  
composition of the seed. Such oil preferably has from about  
74 to about 90% oleic acid, more preferably from about 80 to  
20 about 90% oleic acid. In some embodiments, oil obtained  
from seeds produced by plants of the invention may have from  
about 2.0% to about 5.0% saturated fatty acids, based on  
total fatty acid composition of the seeds. In some  
embodiments, oil obtained from seeds of the invention may  
25 have from about 1.0% to about 14.0% linoleic acid, or from  
about 0.5% to about 10.0%  $\alpha$ -linolenic acid.

Oil composition typically is analyzed by crushing  
and extracting fatty acids from bulk seed samples (e.g., 10  
seeds). Fatty acid triglycerides in the seed are hydrolyzed  
30 and converted to fatty acid methyl esters. Those seeds  
having an altered fatty acid composition may be identified  
by techniques known to the skilled artisan, e.g., gas-liquid  
chromatography (GLC) analysis of a bulked seed sample or of

5 a single half-seed. Half-seed analysis is well known in the art to be useful because the viability of the embryo is maintained and thus those seeds having a desired fatty acid profile may be planted to from the next generation.

10 However, half-seed analysis is also known to be an inaccurate representation of genotype of the seed being analyzed. Bulk seed analysis typically yields a more accurate representation of the fatty acid profile of a given genotype. Fatty acid composition can also be determined on larger samples, e.g., oil obtained by pilot plant or commercial scale refining, bleaching and deodorizing of endogenous oil in the seeds.

15 The nucleic acid fragments of the invention can be used as markers in plant genetic mapping and plant breeding programs. Such markers may include restriction fragment length polymorphism (RFLP), random amplification polymorphism detection (RAPD), polymerase chain reaction (PCR) or self-sustained sequence replication (3SR) markers, for example. Marker-assisted breeding techniques may be used to identify and follow a desired fatty acid composition during the breeding process. Marker-assisted breeding techniques may be used in addition to, or as an alternative to, other sorts of identification techniques. An example of marker-assisted breeding is the use of PCR primers that specifically amplify a sequence containing a desired mutation in delta-12 desaturase or delta-15 desaturase.

25 Methods according to the invention are useful in that the resulting plants and plant lines have desirable seed fatty acid compositions as well as superior agronomic properties compared to known lines having altered seed fatty acid composition. Superior agronomic characteristics include, for example, increased seed germination percentage, increased seedling vigor, increased resistance to seedling

09271904-012901  
fungal diseases (damping off, root rot and the like),  
increased yield, and improved standability.

While the invention is susceptible to various  
modifications and alternative forms, certain specific  
5 embodiments thereof are described in the general methods and  
examples set forth below. For example the invention may be  
applied to all *Brassica* species, including *B. rapa*, *B.*  
*juncea*, and *B. hirta*, to produce substantially similar  
results. It should be understood, however, that these  
10 examples are not intended to limit the invention to the  
particular forms disclosed but, instead the invention is to  
cover all modifications, equivalents and alternatives  
falling within the scope of the invention. This includes  
the use of somaclonal variation; physical or chemical  
15 mutagenesis of plant parts; anther, microspore or ovary  
culture followed by chromosome doubling; or self- or cross-  
pollination to transmit the fatty acid trait, alone or in  
combination with other traits, to develop new *Brassica*  
lines.

#### 20 EXAMPLE 1

##### Mutagenesis

Seeds of Westar, a Canadian (*Brassica napus*) spring  
canola variety, were subjected to chemical mutagenesis.  
Westar is a registered Canadian spring variety with canola  
25 quality. The fatty acid composition of field-grown Westar,  
3.9% C<sub>16:0</sub>, 1.9% C<sub>18:0</sub>, 67.5% C<sub>18:1</sub>, 17.6% C<sub>18:2</sub>, 7.4% C<sub>18:3</sub>, <2%  
C<sub>20:1</sub> + C<sub>22:1</sub>, has remained stable under commercial  
production, with  $\pm 10\%$  deviation, since 1982.

Prior to mutagenesis, 30,000 seeds of *B. napus* cv.  
30 Westar seeds were preimbibed in 300-seed lots for two hours  
on wet filter paper to soften the seed coat. The preimbibed  
seeds were placed in 80 mM ethylmethanesulfonate (EMS) for



four hours. Following mutagenesis, the seeds were rinsed three times in distilled water. The seeds were sown in 48-well flats containing Pro-Mix. Sixty-eight percent of the mutagenized seed germinated. The plants were maintained at 25°C/15°C, 14/10 hr day/night conditions in the greenhouse. At flowering, each plant was individually self-pollinated.

M<sub>2</sub> seed from individual plants were individually catalogued and stored, approximately 15,000 M<sub>2</sub> lines was planted in a summer nursery in Carman, Manitoba. The seed from each selfed plant were planted in 3-meter rows with 6-inch row spacing. Westar was planted as the check variety. Selected lines in the field were selfed by bagging the main raceme of each plant. At maturity, the selfed plants were individually harvested and seeds were catalogued and stored to ensure that the source of the seed was known.

Self-pollinated M<sub>3</sub> seed and Westar controls were analyzed in 10-seed bulk samples for fatty acid composition via gas chromatography. Statistical thresholds for each fatty acid component were established using a Z-distribution with a stringency level of 1 in 10,000. Mean and standard deviation values were determined from the non-mutagenized Westar control population in the field. The upper and lower statistical thresholds for each fatty acid were determined from the mean value of the population  $\pm$  the standard deviation, multiplied by the Z-distribution. Based on a population size of 10,000, the confidence interval is 99.99%.

The selected M<sub>3</sub> seeds were planted in the greenhouse along with Westar controls. The seed was sown in 4-inch pots containing Pro-Mix soil and the plants were maintained at 25°C/15°C, 14/10 hr day/night cycle in the greenhouse. At flowering, the terminal raceme was self-pollinated by bagging. At maturity, selfed M<sub>4</sub> seed was individually

harvested from each plant, labelled, and stored to ensure that the source of the seed was known.

The  $M_4$  seed was analyzed in 10-seed bulk samples. Statistical thresholds for each fatty acid component were established from 259 control samples using a Z-distribution of 1 in 800. Selected  $M_4$  lines were planted in a field trial in Carman, Manitoba in 3-meter rows with 6-inch spacing. Ten  $M_4$  plants in each row were bagged for self-pollination. At maturity, the selfed plants were individually harvested and the open pollinated plants in the row were bulk harvested. The  $M_5$  seed from single plant selections was analyzed in 10-seed bulk samples and the bulk row harvest in 50-seed bulk samples.

Selected  $M_5$  lines were planted in the greenhouse along with Westar controls. The seed was grown as previously described. At flowering the terminal raceme was self-pollinated by bagging. At maturity, selfed  $M_6$  seed was individually harvested from each plant and analyzed in 10-seed bulk samples for fatty acid composition.

Selected  $M_6$  lines were entered into field trials in Eastern Idaho. The four trial locations were selected for the wide variability in growing conditions. The locations included Burley, Tetonia, Lamont and Shelley (Table 7). The lines were planted in four 3-meter rows with an 8-inch spacing, each plot was replicated four times. The planting design was determined using a Randomized Complete Block Designed. The commercial cultivar Westar was used as a check cultivar. At maturity the plots were harvested to determine yield. Yield of the entries in the trial was determined by taking the statistical average of the four replications. The Least Significant Difference Test was used to rank the entries in the randomized complete block design.

TABLE 7

Trial Locations for Selected Fatty Acid Mutants

<u>LOCATION</u>	<u>SITE CHARACTERIZATIONS</u>
BURLEY	Irrigated. Long season. High temperatures during flowering.
5 TETONIA	Dryland. Short season. Cool temperatures.
LAMONT	Dryland. Short season. Cool temperatures.
SHELLEY	Irrigated. Medium season. High temperatures during flowering.

To determine the fatty acid profile of entries, plants in each plot were bagged for self-pollination. The  
10 M<sub>1</sub> seed from single plants was analyzed for fatty acids in ten-seed bulk samples.

To determine the genetic relationships of the selected fatty acid mutants crosses were made. Flowers of M<sub>6</sub> or later generation mutations were used in crossing. F<sub>1</sub>  
15 seed was harvested and analyzed for fatty acid composition to determine the mode of gene action. The F<sub>1</sub> progeny were planted in the greenhouse. The resulting plants were self-pollinated, the F<sub>2</sub> seed harvested and analyzed for fatty acid composition for allelism studies. The F<sub>2</sub> seed and  
20 parent line seed was planted in the greenhouse, individual plants were self-pollinated. The F<sub>3</sub> seed of individual plants was tested for fatty acid composition using 10-seed bulk samples as described previously.

In the analysis of some genetic relationships  
25 dihaploid populations were made from the microspores of the F<sub>1</sub> hybrids. Self-pollinated seed from dihaploid plants were analyzed for fatty acid analysis using methods described previously.

For chemical analysis, 10-seed bulk samples were  
30 hand ground with a glass rod in a 15-mL polypropylene tube

and extracted in 1.2 mL 0.25 N KOH in 1:1 ether/methanol. The sample was vortexed for 30 sec. and heated for 60 sec. in a 60°C water bath. Four mL of saturated NaCl and 2.4 mL of iso-octane were added, and the mixture was vortexed again. After phase separation, 600 µL of the upper organic phase were pipetted into individual vials and stored under nitrogen at -5°C. One µL samples were injected into a Supelco SP-2330 fused silica capillary column (0.25 mm ID, 30 M length, 0.20 µm df).

10 The gas chromatograph was set at 180°C for 5.5 minutes, then programmed for a 2°C/minute increase to 212°C, and held at this temperature for 1.5 minutes. Total run time was 23 minutes. Chromatography settings were: Column head pressure - 15 psi, Column flow (He) - 0.7 mL/min.,  
15 Auxiliary and Column flow - 33 mL/min., Hydrogen flow - 33 mL/min., Air flow - 400 mL/min., Injector temperature - 250°C, Detector temperature - 300°C, Split vent - 1/15.

Table 8 describes the upper and lower statistical thresholds for each fatty acid of interest.

**TABLE 8**

Statistical Thresholds for Specific Fatty Acids  
Derived from Control Westar Plantings

5	Genotype	Percent Fatty Acids				
		C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>
	M <sub>3</sub> Generation (1 in 10,000 rejection rate)					
	Lower	3.3	1.4	--	13.2	5.3
	Upper	4.3	2.5	71.0	21.6	9.9
10	M <sub>4</sub> Generation (1 in 800 rejection rate)					
	Lower	3.6	0.8	--	12.2	3.2
	Upper	6.3	3.1	76.0	32.4	9.9
	M <sub>5</sub> Generation (1 in 755 rejection rate)					
	Lower	2.7	0.9	--	9.6	2.6
15	Upper	5.7	2.7	80.3	26.7	9.6

\*Sats=Total Saturate Content

**EXAMPLE 2**

High Oleic Acid Canola Lines

In the studies of Example 1, at the M<sub>3</sub> generation, 20 31 lines exceeded the upper statistical threshold for oleic acid ( $\geq 71.0\%$ ). Line W7608.3 had 71.2% oleic acid. At the M<sub>4</sub> generation, its selfed progeny (W7608.3.5, since designated A129.5) continued to exceed the upper statistical threshold for C<sub>18:1</sub> with 78.8% oleic acid. M<sub>5</sub> seed of five 25 self-pollinated plants of line A129.5 (ATCC 40811) averaged 75.0% oleic acid. A single plant selection, A129.5.3 had 75.6% oleic acid. The fatty acid composition of this high oleic acid mutant, which was stable under both field and greenhouse conditions to the M<sub>5</sub> generation, is summarized in

Table 9. This line also stably maintained its mutant fatty acid composition to the M<sub>7</sub> generation in field trials in multiple locations. Over all locations the self-pollinated plants (A129) averaged 78.3% oleic acid. The fatty acid composition of the A129 for each Idaho trial location are summarized in Table 10. In multiple location replicated yield trials, A129 was not significantly different in yield from the parent cultivar Westar.

The canola oil of A129, after commercial processing, was found to have superior oxidative stability compared to Westar when measured by the Accelerated Oxygen Method (AOM), American Oil Chemists' Society Official Method Cd 12-57 for fat stability; Active Oxygen Method (revised 1989). The AOM of Westar was 18 AOM hours and for A129 was 30 AOM hours.

**TABLE 9**

Fatty Acid Composition of a High  
Oleic Acid Canola Line Produced by Seed Mutagenesis

		Percent Fatty Acids				
Genotype	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats
Westar	3.9	1.9	67.5	17.6	7.4	7.0
W7608.3 (M <sub>3</sub> )	3.9	2.4	71.2	12.7	6.1	7.6
W7608.3.5 (M <sub>4</sub> )	3.9	2.0	78.8	7.7	3.9	7.3
A129.5.3 (M <sub>5</sub> )	3.8	2.3	75.6	9.5	4.9	7.6

Sats=Total Saturate Content

**TABLE 10**

**Fatty Acid Composition of a Mutant High Oleic Acid Line at Different Field Locations in Idaho**

Percent Fatty Acids						
Location	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats
Burley	3.3	2.1	77.5	8.1	6.0	6.5
Tetonia	3.5	3.4	77.8	6.5	4.7	8.5
Lamont	3.4	1.9	77.8	7.4	6.5	6.3
Shelley	3.3	2.6	80.0	5.7	4.5	7.7
Sats=Total Saturate Content						

The genetic relationship of the high oleic acid mutation A129 to other oleic desaturases was demonstrated in crosses made to commercial canola cultivars and a low linolenic acid mutation. A129 was crossed to the commercial cultivar Global (C<sub>16:0</sub> - 4.5%, C<sub>18:0</sub> - 1.5%, C<sub>18:1</sub> - 62.9%, C<sub>18:2</sub> - 20.0%, C<sub>18:3</sub> - 7.3%). Approximately 200 F<sub>2</sub> individuals were analyzed for fatty acid composition. The results are summarized in Table 11. The segregation fit 1:2:1 ratio suggesting a single co-dominant gene controlled the inheritance of the high oleic acid phenotype.

**TABLE 11**

**Genetic Studies of A129 X Global**

Frequency			
Genotype	C <sub>18:0</sub> Content (%)	Observed	Expected
od-od-	77.3	43	47
od-od+	71.7	106	94
od+od+	66.1	49	47

A cross between A129 and IMC 01, a low linolenic acid variety (C<sub>16:0</sub> - 4.1%, C<sub>18:0</sub> - 1.9%, C<sub>18:1</sub> - 66.4%, C<sub>18:2</sub> - 18.1%, C<sub>18:3</sub> - 5.7%), was made to determine the inheritance of the oleic acid desaturase and linoleic acid desaturase.

In the F<sub>1</sub> hybrids both the oleic acid and linoleic acid desaturase genes approached the mid-parent values indicating a co-dominant gene actions. Fatty acid analysis of the F<sub>2</sub> individuals confirmed a 1:2:1:2:4:2:1:2:1 segregation of two independent, co-dominant genes (Table 12). A line was selected from the cross of A129 and IMC01 and designated as IMC130 (ATCC deposit no. 75446) as described in U.S. Patent Application No. 08/425,108, incorporated herein by reference.

**TABLE 12**

Genetic Studies of A129 X IMC 01

	Genotype	Ratio	Frequency	
			Observed	Expected
	od-od-ld-ld-	1	11	12
15	od-od-ld-ld+	2	30	24
	od-od-ld+ld+	1	10	12
	od-od+ld-ld-	2	25	24
	od-od+ld-ld+	4	54	47
	od-od+ld+ld+	2	18	24
20	od+od+ld-ld-	1	7	12
	od+od+ld-ld+	2	25	24
	od+od+ld+ld+	1	8	12

An additional high oleic acid line, designated A128.3, was also produced by the disclosed method. A 50-seed bulk analysis of this line showed the following fatty acid composition: C<sub>16:0</sub> - 3.5%, C<sub>18:0</sub> - 1.8%, C<sub>18:1</sub> - 77.3%, C<sub>18:2</sub> - 9.0%, C<sub>18:3</sub> - 5.6%, FDA Sats - 5.3%, Total Sats - 6.4%.

This line also stably maintained its mutant fatty acid composition to the M<sub>7</sub> generation. In multiple locations replicated yield trials, A128 was not significantly different in yield from the parent cultivar Westar.

A129 was crossed to A128.3 for allelism studies. Fatty acid composition of the F<sub>2</sub> seed showed the two lines to be allelic. The mutational events in A129 and A128.3 although different in origin were in the same gene.



097704-01201

An additional high oleic acid line, designated M3028.-10 (ATCC 75026), was also produced by the disclosed method in Example 1. A 10-seed bulk analysis of this line showed the following fatty acid composition: C<sub>16:0</sub> - 3.5%, C<sub>18:0</sub> - 1.8%, C<sub>18:1</sub> - 77.3%, C<sub>18:2</sub> - 9.0%, C<sub>18:3</sub> - 5.6%, FDA Saturates - 5.3%, Total Saturates - 6.4%. In a single location replicated yield trial M3028.10 was not significantly different in yield from the parent cultivar Westar.

### EXAMPLE 3

#### Low Linoleic Acid Canola

In the studies of Example 1, at the M<sub>3</sub> generation, 80 lines exceeded the lower statistical threshold for linoleic acid ( $\leq 13.2\%$ ). Line W12638.8 had 9.4% linoleic acid. At the M<sub>4</sub> and M<sub>5</sub> generations, its selfed progenies [W12638.8, since designated A133.1 (ATCC 40812)] continued to exceed the statistical threshold for low C<sub>18:2</sub> with linoleic acid levels of 10.2% and 8.4%, respectively. The fatty acid composition of this low linoleic acid mutant, which was stable to the M<sub>5</sub> generation under both field and greenhouse conditions, is summarized in Table 13. In multiple location replicated yield trials, A133 was not significantly different in yield from the parent cultivar Westar. An additional low linoleic acid line, designated M3062.8 (ATCC 75025), was also produced by the disclosed method. A 10-seed bulk analysis of this line showed the following fatty acid composition: C<sub>16:0</sub> - 3.8%, C<sub>18:0</sub> - 2.3%, C<sub>18:1</sub> - 77.1%, C<sub>18:2</sub> - 8.9%, C<sub>18:3</sub> - 4.3%, FDA Sats-6.1%. This line has also stably maintained its mutant fatty acid composition in the field and greenhouse.

TABLE 13

Fatty Acid Composition of a Low  
Linoleic Acid Canola Line Produced by Seed Mutagenesis

		Percent Fatty Acids				
	Genotype	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	Sats <sup>b</sup>
5	Westar	3.9	1.9	67.5	17.6	7.4
	W12638.8 (M <sub>3</sub> )	3.9	2.3	75.0	9.4	6.1
	W12638.8.1 (M <sub>4</sub> )	4.1	1.7	74.6	10.2	5.9
10	A133.1.8 (M <sub>5</sub> )	3.8	2.0	77.7	8.4	5.0
						7.0

<sup>a</sup>Letter and numbers up to second decimal point indicate the plant line. Number after second decimal point indicates an individual plant.

<sup>b</sup>Sats=Total Saturate Content

## EXAMPLE 4

Low Linolenic and Linoleic Acid Canola

In the studies of Example 1, at the M<sub>3</sub> generation, 57 lines exceeded the lower statistical threshold for linolenic acid ( $\leq 5.3\%$ ). Line W14749.8 had 5.3% linolenic acid and 15.0% linoleic acid. At the M<sub>4</sub> and M<sub>5</sub> generations, its selfed progenies [W14749.8, since designated M3032 (ATCC 75021)] continued to exceed the statistical threshold for low C<sub>18:3</sub> with linolenic acid levels of 2.7% and 2.3%, respectively, and for a low sum of linolenic and linoleic acids with totals of 11.8% and 12.5% respectively. The fatty acid composition of this low linolenic acid plus linoleic acid mutant, which was stable to the M<sub>5</sub> generation under both field and greenhouse conditions, is summarized in Table 14. In a single location replicated yield trial M3032

was not significantly different in yield from the parent cultivar (Westar).

TABLE 14

Fatty Acid Composition of a Low  
Linolenic Acid Canola Line Produced by Seed Mutagenesis

Genotype	Percent Fatty Acids					Sats
	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	
Westar	3.9	1.9	67.5	17.6	7.4	7.0
W14749.8 (M <sub>3</sub> )	4.0	2.5	69.4	15.0	5.3	6.5
M3032.8 (M <sub>4</sub> )	3.9	2.4	77.9	9.1	2.7	6.4
M3032.1 (M <sub>5</sub> )	3.5	2.8	80.0	10.2	2.3	6.5

Sats=Total Saturate Content

EXAMPLE 5

Canola Lines Q508 and Q4275

Seeds of the *B. napus* line IMC-129 were mutagenized with methyl N-nitrosoguanidine (MNNG). The MNNG treatment consisted of three parts: pre-soak, mutagen application, and wash. A 0.05M Sorenson's phosphate buffer was used to maintain pre-soak and mutagen treatment pH at 6.1. Two hundred seeds were treated at one time on filter paper (Whatman #3M) in a petri dish (100mm x 15mm). The seeds were pre-soaked in 15 mls of 0.05M Sorenson's buffer, pH 6.1, under continued agitation for two hours. At the end of the pre-soak period, the buffer was removed from the plate.

A 10mM concentration of MNNG in 0.05M Sorenson's buffer, pH 6.1, was prepared prior to use. Fifteen ml of 10m MNNG was added to the seeds in each plate. The seeds were incubated at 22°C±3°C in the dark under constant

agitation for four (4) hours. At the end of the incubation period, the mutagen solution was removed.

The seeds were washed with three changes of distilled water at 10 minute intervals. The fourth wash was for thirty minutes. This treatment regime produced an LD60 population.

Treated seeds were planted in standard greenhouse potting soil and placed into an environmentally controlled greenhouse. The plants were grown under sixteen hours of light. At flowering, the racemes were bagged to produce selfed seed. At maturity, the M2 seed was harvested. Each M2 line was given an identifying number. The entire MNNG-treated seed population was designated as the Q series.

Harvested M2 seeds were planted in the greenhouse. The growth conditions were maintained as previously described. The racemes were bagged at flowering for selfing. At maturity, the selfed M3 seed was harvested and analyzed for fatty acid composition. For each M3 seed line, approximately 10-15 seeds were analyzed in bulk as described in Example 1.

High oleic-low linoleic M3 lines were selected from the M3 population using a cutoff of >82% oleic acid and <5.0% linoleic. From the first 1600 M3 lines screened for fatty acid composition, Q508 was identified. The Q508 M3 generation was advanced to the M4 generation in the greenhouse. Table 15 shows the fatty acid composition of Q508 and IMC 129. The M4 selfed seed maintained the selected high oleic-low linoleic acid phenotype (Table 16).

**TABLE 15**

Fatty Acid Composition of A129 and High  
Oleic Acid M3 Mutant Q508

Line #	16:0	18:0	18:1	18:2	18:3
5 A129 <sup>*</sup>	4.0	2.4	77.7	7.8	4.2
Q508	3.9	2.1	84.9	2.4	2.9

<sup>\*</sup>Fatty acid composition of A129 is the average of  
50 self-pollinated plants grown with the M3 population

10 M<sub>4</sub> generation Q508 plants had poor agronomic  
qualities in the field compared to Westar. Typical plants  
were slow growing relative to Westar, lacked early  
vegetative vigor, were short in stature, tended to be  
chlorotic and had short pods. The yield of Q508 was very  
low compared to Westar.

15 The M<sub>4</sub> generation Q508 plants in the greenhouse  
tended to be reduced in vigor compared to Westar. However,  
Q508 yields in the greenhouse were greater than Q508 yields  
in the field.

**TABLE 16**

Fatty Acid Composition of Seed Oil  
from Greenhouse-Grown Q508, IMC 129 and Westar.

Line	16:0	18:0	18:1	18:2	18:3	FDA Sats
IMC 129 <sup>a</sup>	4.0	2.4	77.7	7.8	4.2	6.4
25 Westar <sup>b</sup>	3.9	1.9	67.5	17.6	7.4	>5.8
Q508 <sup>c</sup>	3.9	2.1	84.9	2.4	2.9	6.0

<sup>a</sup>Average of 50 self-pollinated plants

<sup>b</sup>Data from Example 1

<sup>c</sup>Average of 50 self-pollinated plants

Nine other M4 high-oleic low-linoleic lines were also identified: Q3603, Q3733, Q4249, Q6284, Q6601, Q6761, Q7415, Q4275, and Q6676. Some of these lines had good agronomic characteristics and an elevated oleic acid level in seeds of about 80% to about 84%.

Q4275 was crossed to the variety Cyclone. After selfing for seven generations, mature seed was harvested from 93GS34-179, a progeny line of the Q4275 Cyclone cross. Referring to Table 17, fatty acid composition of a bulk seed sample shows that 93GS34 retained the seed fatty acid composition of Q4275. 93GS34-179 also maintained agronomically desirable characteristics.

After more than seven generations of selfing of Q4275, plants of Q4275, IMC 129 and 93GS34 were field grown during the summer season. The selections were tested in 4 replicated plots (5 feet X 20 feet) in a randomized block design. Plants were open pollinated. No selfed seed was produced. Each plot was harvested at maturity, and a sample of the bulk harvested seed from each line was analyzed for fatty acid composition as described above. The fatty acid compositions of the selected lines are shown in Table 17.

**TABLE 17**

Fatty Acid Composition of  
Field Grown IMC 129, Q4275 and 93GS34 Seeds

Line	Fatty Acid Composition (%)					
	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	FDA Sats.
IMC 129	3.3	2.4	76.7	8.7	5.2	5.7
Q4275	3.7	3.1	82.1	4.0	3.5	6.8
93GS34-179	2.6	2.7	85.0	2.8	3.3	5.3

0971904-02904  
The results shown in Table 17 show that Q4275 maintained the selected high oleic - low linoleic acid phenotype under field conditions. The agronomic characteristics of Q4275 plants were superior to those of Q508.

M<sub>4</sub> generation Q508 plants were crossed to a dihaploid selection of Westar, with Westar serving as the female parent. The resulting F<sub>1</sub> seed was termed the 92EF population. About 126 F<sub>1</sub> individuals that appeared to have better agronomic characteristics than the Q508 parent were selected for selfing. A portion of the F<sub>2</sub> seed from such individuals was replanted in the field. Each F<sub>2</sub> plant was selfed and a portion of the resulting F<sub>3</sub> seed was analyzed for fatty acid composition. The content of oleic acid in F<sub>3</sub> seed ranged from 59 to 79%. No high oleic (>80%) individuals were recovered with good agronomic type.

A portion of the F<sub>2</sub> seed of the 92EF population was planted in the greenhouse to analyze the genetics of the Q508 line. F<sub>3</sub> seed was analyzed from 380 F<sub>2</sub> individuals. The C<sub>18:1</sub> levels of F<sub>3</sub> seed from the greenhouse experiment is depicted in Figure 1. The data were tested against the hypothesis that Q508 contains two mutant genes that are semi-dominant and additive: the original IMC 129 mutation as well as one additional mutation. The hypothesis also assumes that homozygous Q508 has greater than 85% oleic acid and homozygous Westar has 62-67% oleic acid. The possible genotypes at each gene in a cross of Q508 by Westar may be designated as:

AA = Westar Fad2<sup>a</sup>

BB = Westar Fad2<sup>b</sup>

aa = Q508 Fad2<sup>a-</sup>

bb = Q508 Fad2<sup>b-</sup>

Assuming independent segregation, a 1:4:6:4:1 ratio of phenotypes is expected. The phenotypes of heterozygous plants are assumed to be indistinguishable and, thus, the data were tested for fit to a 1:14:1 ratio of homozygous Westar: heterozygous plants: homozygous Q508.

Phenotypic Ratio	# of Westar Alleles	Genotype
1	4	AABB (Westar)
4	3	AABb, AaBB, AABb, AaBB
6	2	AaBb, AAbb, AaBb, AaBb, aaBB, AaBb
4	1	Aabb, aaBb, Aabb, aaBb
1	0	aabb (Q508)

Using Chi-square analysis, the oleic acid data fit a 1:14:1 ratio. It was concluded that Q508 differs from Westar by two major genes that are semi-dominant and additive and that segregate independently. By comparison, the genotype of IMC 129 is aaBB.

The fatty acid composition of representative F3 individuals having greater than 85% oleic acid in seed oil is shown in Table 18. The levels of saturated fatty acids are seen to be decreased in such plants, compared to Westar.



**TABLE 18**

92EF F, Individuals with >85% C<sub>18:1</sub> in Seed Oil

F3 Plant Identifier	Fatty Acid Composition (%)					
	C16:0	C18:0	C18:1	C18:2	C18:3	FDASA
+38068	3.401	1.582	85.452	2.134	3.615	4.983
+38156	3.388	1.379	85.434	2.143	3.701	4.767
+38171	3.588	1.511	85.289	2.367	3.425	5.099
+38181	3.75	1.16	85.312	2.968	3.819	4.977
+38182	3.529	0.985	85.905	2.614	3.926	4.56
+38191	3.364	1.039	85.737	2.869	4.039	4.459
+38196	3.557	1.182	85.054	2.962	4.252	4.739
+38202	3.554	1.105	86.091	2.651	3.721	4.713
+38220	3.093	1.16	86.421	1.931	3.514	4.314
+38236	3.308	1.349	85.425	2.37	3.605	4.718
+38408	3.617	1.607	85.34	2.33	3.562	5.224
+38427	3.494	1.454	85.924	2.206	3.289	4.948
+38533	3.64	1.319	85.962	2.715	3.516	4.959

**EXAMPLE 6**

Leaf and Root Fatty Acid Profiles of Canola Lines IMC-129, Q508, and Westar

Plants of Q508, IMC 129 and Westar were grown in the greenhouse. Mature leaves, primary expanding leaves, petioles and roots were harvested at the 6-8 leaf stage,

09771904-012901  
T062F0-406T2260  
frozen in liquid nitrogen and stored at  $-70^{\circ}\text{C}$ . Lipid extracts were analyzed by GLC as described in Example 1. The fatty acid profile data are shown in Table 19.

5       The data in Table 19 indicate that total leaf lipids in Q508 are higher in  $\text{C}_{18:1}$  content than the  $\text{C}_{18:2}$  plus  $\text{C}_{18:3}$  content. The reverse is true for Westar and IMC 129. The difference in total leaf lipids between Q508 and IMC 129 is consistent with the hypothesis that a second Fad2 gene is mutated in Q508.

10       The  $\text{C}_{16:3}$  content in the total lipid fraction was about the same for all three lines, suggesting that the plastid FadC gene product was not affected by the Q508 mutations. To confirm that the FadC gene was not mutated, chloroplast lipids were separated and analyzed. No changes  
15   in chloroplast  $\text{C}_{16:1}$ ,  $\text{C}_{16:2}$  or  $\text{C}_{16:3}$  fatty acids were detected in the three lines. The similarity in plastid leaf lipids among Q508, Westar and IMC 129 is consistent with the hypothesis that the second mutation in Q508 affects a microsomal Fad2 gene and not a plastid FadC gene.

TABLE 19

	MATURE LEAF			EXPANDING LEAF			PETIOLE			ROOT		
	West.	129	3Q508	West.	129	3Q508	West.	129	3Q508	West.	129	3Q508
16:0	12.1	11.9	10.1	16.4	16.1	11.3	21.7	23.5	11.9	21.1	21.9	12.0
16:1	0.8	0.6	1.1	0.7	0.6	1.1	1.0	1.3	1.4	-	-	-
16:2	2.3	2.2	2.0	2.8	3.1	2.8	1.8	2.2	1.8	-	-	-
16:3	14.7	15.0	14.3	6.3	5.4	6.9	5.7	4.6	5.7	-	-	-
18:0	2.2	1.6	1.2	2.5	2.8	1.5	3.7	4.0	1.6	3.6	2.9	2.5
18:1	2.8	4.9	16.7	3.8	8.3	38.0	4.9	12.9	46.9	3.5	6.1	68.8
18:2	12.6	11.5	6.8	13.3	13.8	4.9	20.7	18.3	5.2	28.0	30.4	4.4
18:3	50.6	50.3	46.0	54.2	50.0	33.5	40.4	33.2	25.3	43.8	38.7	12.3

## EXAMPLE 7

Sequences of Mutant and Wild-Type Delta-12 Fatty Acid Desaturases from *B. napus*

Primers specific for the FAD2 structural gene were used to clone the entire open reading frame (ORF) of the D and F delta-12 desaturase genes by reverse transcriptase polymerase chain reaction (RT-PCR). RNA from seeds of IMC 129, Q508 and Westar plants was isolated by standard methods and was used as template. The RT-amplified fragments were used for nucleotide sequence determination. The DNA sequence of each gene from each line was determined from both strands by standard dideoxy sequencing methods.

Sequence analysis revealed a G to A transversion at nucleotide 316 (from the translation initiation codon) of the D gene in both IMC 129 and Q508, compared to the sequence of Westar. The transversion changes the codon at this position from GAG to AAG and results in a non-conservative substitution of glutamic acid, an acidic residue, for lysine a basic residue. The presence of the

same mutation in both lines was expected since the Q508 line was derived from IMC 129. The same base change was also detected in Q508 and IMC 129 when RNA from leaf tissue was used as template.

5           The G to A mutation at nucleotide 316 was confirmed by sequencing several independent clones containing fragments amplified directly from genomic DNA of IMC 129 and Westar. These results eliminated the possibility of a rare mutation introduced during reverse transcription and PCR in  
10 the RT-PCR protocol. It was concluded that the IMC 129 mutant is due to a single base transversion at nucleotide 316 in the coding region of the D gene of rapeseed microsomal delta 12-desaturase.

          A single base transition from T to A at nucleotide  
15 515 of the F gene was detected in Q508 compared to the Westar sequence. The mutation changes the codon at this position from CTC to CAC, resulting in the non-conservative substitution of a ~~non-polar residue, leucine, for a polar residue, histidine~~<sup>A</sup>, in the resulting gene product. No  
20 mutations were found in the F gene sequence of IMC 129 compared to the F gene sequence of Westar.

          These data support the conclusion that a mutation in a delta-12 desaturase gene sequence results in alterations in the fatty acid profile of plants containing such a  
25 mutated gene. Moreover, the data show that when a plant line or species contains two delta-12 desaturase loci, the fatty acid profile of an individual having two mutated loci differs from the fatty acid profile of an individual having one mutated locus.

30           The mutation in the D gene of IMC 129 and Q508 mapped to a region having a conserved amino acid motif (His-Xaa-Xaa-Xaa-His) found in cloned delta-12 and delta-15 membrane bound-desaturases (Table 20).

TABLE 20

Alignment of Amino Acid Sequences  
of Cloned Canola Membrane Bound-Desaturases

Desaturase Gene	Sequence <sup>a</sup>	Position
Canola-fad2-D (mutant)	AHKCGH	109-114
Canola-Fad2-D	AHECGH	109-114
Canola-Fad2-F	AHECGH	109-114
Canola-FadC	<u>GHDCAH</u>	170-175
Canola-fad3 (mutant)	<u>GHKCGH</u>	94-99
Canola-Fad3	<u>GHDCGH</u>	94-99
Canola-FadD	<u>GHDCGH</u>	125-130

(FadD = Plastid delta 15, Fad3 = Microsomal delta-15),  
(FadC = Plastid delta-12, Fad2 = Microsomal delta-12)

<sup>a</sup> One letter amino acid code; conservative substitutions are underlined; non-conservative substitutions are in bold.

EXAMPLE 8

Transcription and Translation of Microsomal Delta-12  
Fatty Acid Desaturases

Transcription in vivo was analyzed by RT-PCR analysis of stage II and stage III developing seeds and leaf tissue. The primers used to specifically amplify delta-12 desaturase F gene RNA from the indicated tissues were sense primer 5'-GGATATGATGATGGT<sup>(Seq ID NO: 19)</sup>GAAAGA-3' and antisense primer 5'-TCTTTCACCATCATCATATCC-3'<sup>(Seq ID NO: 20)</sup>. The primers used to specifically amplify delta-12 desaturase D gene RNA from the indicated tissues were sense primer 5'-GTTATGAAGCAAAGAAGAAAC-3'<sup>(Seq ID NO: 21)</sup> and antisense primer 5'-GTTTCTTCTTTGCTTCATAAC-3'<sup>(Seq ID NO: 22)</sup>. The results indicated that mRNA of both the D and F gene was expressed

Sub  
C100  
in seed and leaf tissues of IMC 129, Q508 and wild type Westar plants.

In vitro transcription and translation analysis showed that a peptide of about 46 kD was made. This is the expected size of both the D gene product and the F gene product, based on sum of the deduced amino acid sequence of each gene and the cotranslational addition of a microsomal membrane peptide.

These results rule out the possibility that non-sense or frameshift mutations, resulting in a truncated polypeptide gene product, are present in either the mutant D gene or the mutant F gene. The data, in conjunction with the data of Example 7, support the conclusion that the mutations in Q508 and IMC 129 are in delta-12 fatty acid desaturase structural genes encoding desaturase enzymes, rather than in regulatory genes.

#### EXAMPLE 9

##### Development of Gene-Specific PCR Markers

Based on the single base change in the mutant D gene of IMC 129 described in above, two 5' PCR primers were designed. The nucleotide sequence of the primers differed only in the base (G for Westar and A for IMC 129) at the 3' end. The primers allow one to distinguish between mutant fad2-D and wild-type Fad2-D alleles in a DNA-based PCR assay. Since there is only a single base difference in the 5' PCR primers, the PCR assay is very sensitive to the PCR conditions such as annealing temperature, cycle number, amount, and purity of DNA templates used. Assay conditions have been established that distinguish between the mutant gene and the wild type gene using genomic DNA from IMC 129 and wild type plants as templates. Conditions may be further optimized by varying PCR parameters, particularly

with variable crude DNA samples. A PCR assay distinguishing the single base mutation in IMC 129 from the wild type gene along with fatty acid composition analysis provides a means to simplify segregation and selection analysis of genetic crosses involving plants having a delta-12 fatty acid desaturase mutation.

#### EXAMPLE 10

##### Transformation with Mutant and Wild Type Fad3 Genes

*B. napus* cultivar Westar was transformed with mutant and wild type Fad3 genes to demonstrate that the mutant Fad3 gene for canola cytoplasmic linoleic desaturase delta-15 desaturase is nonfunctional. Transformation and regeneration were performed using disarmed *Agrobacterium tumefaciens* essentially following the procedure described in WO 94/11516.

Two disarmed *Agrobacterium* strains were engineered, each containing a Ti plasmid having the appropriate gene linked to a seed-specific promoter and a corresponding termination sequence. The first plasmid, pIMC110, was prepared by inserting into a disarmed Ti vector the full length wild type Fad3 gene in sense orientation (nucleotides 208 to 1336 of SEQ ID 6 in WO 93/11245), flanked by a napin promoter sequence positioned 5' to the Fad3 gene and a napin termination sequence positioned 3' to the Fad3 gene. The rapeseed napin promoter is described in EP 0255378.

The second plasmid, pIMC205, was prepared by inserting a mutated Fad3 gene in sense orientation into a disarmed Ti vector. The mutant sequence contained mutations at nucleotides 411 and 413 of the microsomal Fad3 gene described in WO93/11245, thus changing the sequence for codon 96 from GAC to AAG. The amino acid at codon 96 of the gene product was thereby changed from aspartic acid to lysine. See Table 20. A bean (*Phaseolus vulgaris*)

Sub  
C8  
cond

phaseolin (7S seed storage protein) promoter fragment of 495 base pairs, starting with 5'-TGGTCTTTTGGT-3', was placed 5' to the mutant Fad3 gene and a phaseolin termination sequence was placed 3' to the mutant Fad3 gene. The phaseolin  
5 sequence is described in Doyle et al., (1986) J. Biol. Chem. 261:9228-9238) and Slightom et al., (1983) Proc. Natl. Acad. Sci. USA 80:1897-1901.

The appropriate plasmids were engineered and transferred separately to Agrobacterium strain LEA4404.  
10 Each engineered strain was used to infect 5 mm segments of hypocotyl explants from Westar seeds by cocultivation. Infected hypocotyls were transferred to callus medium and, subsequently, to regeneration medium. Once discernable  
15 stems formed from the callus, shoots were excised and transferred to elongation medium. The elongated shoots were cut, dipped in Rootone™, rooted on an agar medium and transplanted to potting soil to obtain fertile T1 plants. T2 seeds were obtained by selfing the resulting T1 plants.

Fatty acid analysis of T2 seeds was carried out as  
20 described above. The results are summarized in Table 21. Of the 40 transformants obtained using the pIMC110 plasmid, 17 plants demonstrated wild type fatty acid profiles and 16 demonstrated overexpression. A proportion of the transformants are expected to display an overexpression  
25 phenotype when a functioning gene is transformed in sense orientation into plants.

Of the 307 transformed plants having the pIMC205 gene, none exhibited a fatty acid composition indicative of overexpression. This result indicates that the mutant fad3  
30 gene product is non-functional, since some of the transformants would have exhibited an overexpression phenotype if the gene product were functional.



**TABLE 21**

Overexpression and Co-suppression Events in  
Westar Populations Transformed with pIMC205 or pIMC110.

Construct	Number of Transformants	$\alpha$ -Linolenic Acid Range (%)	Overexpression Events (>10% linolenic)	Co-Suppression Events (<4.0% linolenic)	Wild Type Events
pIMC110	40	2.4 - 20.6	16	7	17
pIMC205	307	4.6 - 10.4	0	0	307

Fatty acid compositions of representative transformed plants are presented in Table 22. Lines 652-09 and 663-40 are representative of plants containing pIMC110 and exhibiting an overexpression and a co-suppression phenotype, respectively. Line 205-284 is representative of plants containing pIMC205 and having the mutant fad3 gene.

**TABLE 22**

Fatty Acid Composition of T2 Seed  
From Westar Transformed With pIMC205 or pIMC110.

Line	Fatty Acid Composition (%)				
	C16:0	C18:0	C18:1	C18:2	C18:3
652-09 pIMC110 overexpression	4.7	3.3	65.6	8.1	14.8
663-40 pIMC110 co-suppression	4.9	2.1	62.5	23.2	3.6
205-284 pIMC205	3.7	1.8	68.8	15.9	6.7

**EXAMPLE 11**

Sequences of Wild Type and Mutant Fad2-D and Fad2-F

High molecular weight genomic DNA was isolated from leaves of Q4275 plants (Example 5). This DNA was used as template for amplification of Fad2-D and Fad2-F genes by polymerase chain reaction (PCR). PCR amplifications were carried out in a total volume of 100  $\mu$ l and contained 0.3  $\mu$ g genomic DNA, 200  $\mu$ M deoxyribonucleoside triphosphates, 3 mM

MgSO<sub>4</sub>, 1-2 Units DNA polymerase and 1X Buffer (supplied by the DNA polymerase manufacturer). Cycle conditions were: 1 cycle for 1 min at 95°C, followed by 30 cycles of 1 min at 94°C, 2 min at 55°C and 3 min at 73°C.

5 The Fad2-D gene was amplified once using Elongase® (Gibco-BRL). PCR primers were: (SEQ ID NO: 23)  
CAUCAUCAUCAUCTTCTTCGTAGGGTTCATCG and (SEQ ID NO: 24)  
CUACUACUACUATCATAGAAGAGAAAGGTTTCAG<sub>^</sub> for the 5' and 3' ends of the gene, respectively.

10 The Fad2-F gene was independently amplified 4 times, twice with Elongase® and twice with Taq polymerase (Boehringer Mannheim). The PCR primers used were: (SEQ ID NO: 25)  
5' CAUCAUCAUCAUATGGGTGCACGTGGAAGAA3' and (SEQ ID NO: 26)  
5' CUACUACUACUATCTTTCACCATCATATCC3' for the 5' and 3' ends of the gene, respectively.

15 Amplified DNA products were resolved on an agarose gel, purified by JetSorb® and then annealed into pAMP1 (Gibco-BRL) via the (CAU)<sub>4</sub> and (CUA)<sub>4</sub> sequences at the ends of the primers, and transformed into *E. coli* DH5α.

20 The Fad2-D and Fad2-F inserts were sequenced on both strands with an ABI PRISM 310 automated sequencer (Perkin-Elmer) following the manufacturer's directions, using synthetic primers, AmpliTaq® DNA polymerase and dye terminator.

25 The Fad2-D gene was found to have an intron upstream of the ATG start codon. As expected, the coding sequence of the gene contained a G to A mutation at nucleotide 316, derived from IMC 129 (Fig. 2).

30 A single base transversion from G to A at nucleotide 908 was detected in the F gene sequence of the Q4275 amplified products, compared to the wild type F gene sequence (Fig. 2). This mutation changes the codon at amino acid 303 from GGA to GAA, resulting in the non-conservative

substitution of a glutamic acid residue for a glycine residue (Table 3 and Fig. 3). Expression of the mutant Q4275 Fad2-F delta-12 desaturase gene in plants alters the fatty acid composition, as described hereinabove.

5           To the extent not already indicated, it will be understood by those of ordinary skill in the art that any one of the various specific embodiments herein described and illustrated may be further modified to incorporate features shown in other of the specific embodiments.

10           The foregoing detailed description has been provided for a better understanding of the invention only and no unnecessary limitation should be understood therefrom as some modifications will be apparent to those skilled in the art without deviating from the spirit and scope of the  
15   appended claims.

0971904-012901